

DIN Malawi Volunteer Medical Pack

This medical pack is issued on behalf of DIN Malawi for volunteers who will be involved in development projects with Development Initiative Network,(DIN Malawi). All volunteers MUST have medical insurance before their trip to Malawi.

Contents About Malawi: The main medical and safety concerns

Before Your Trip: The measures DIN Malawi expects volunteers to take before arriving in Malawi

During Your Stay: How to keep safe while volunteering in Malawi

After Your Visit: Recommendations for preparing for your journey home

About Malawi

Safety: In terms of security, Malawi is perceived as being a very safe place to travel and reside, and Malawians are well known throughout the region for their friendly and welcoming attitudes towards foreigners. The country is roughly the size of England, which makes it relatively easy to travel around, but journeys can be very lengthy nonetheless. The country is landlocked between Mozambique, Tanzania and Zambia, yet since Lake Malawi consumes a fifth of the country's landmass, Malawi is surprisingly green and enjoys a sizeable coastline. Rainy season in the north of the country lasts approximately from November to April, during which time extra precautions should be taken to avoid mosquitoes. There are three main cities: Lilongwe (the Capital), Blantyre and Mzuzu. In most parts of the country tourist facilities are limited and lack of proper infrastructure can cause wild irregularities in electricity, roads, water supply and health care. In general most parts of the country are perceived as safe and secure, mainly because the local communities play such a predominant role in society, especially in rural areas. Street and petty crimes, however, tend to be more common in Lilongwe and secondary cities. You should always watch your bag carefully and carry relatively small amounts of cash. There is a multiparty democracy in Malawi and women have the right to vote. However, female volunteers should be sensitive towards certain cultural differences, especially clothing and should try to dress respectfully and appropriately when passing through towns and villages.

Health: Medical facilities are very basic throughout the country, and outside of the main cities medical resources (doctors, beds, medicines and sanitised equipment) tend to be very limited. Life expectancy stands at approximately 54.8 years - this is due mainly to two factors: HIV and malaria. Around 11% of the adult population has HIV and this number is significantly higher in the rural, northern regions. Malaria claims more than a million lives each year in Africa alone, 90% of them children. It is prevalent in all parts of Malawi, especially during the rainy season (Nov – Apr). In terms of foreigners travelling to Malawi, it has often been documented that there is a risk of catching bilharzia in the otherwise beautiful fresh water of Lake Malawi. Bilharzia is caused by a microscopic worm that lives for part of its life cycle in the human body causing severe stomach upset, exhaustion, and urinary tract infections. It is not a deadly disease and the overall risk is negligible (especially in the north); regardless, travellers should avoid swimming in stagnant water close to human habitation, particularly where there are reeds. Volunteers should also beware of contracting giardia, an infection that affects the small intestine. Giardia is most commonly contracted by drinking contaminated water. Symptoms are very similar to malaria, and it is very easily treated if noticed at an early stage.

Before Your Trip

Vaccinations: All volunteers should visit their GP or a travel health clinic 4 – 8 weeks before departure.

Vaccination Information Hepatitis A Recommended for all volunteers, preferably 4 weeks before departure. A booster should be given 6 – 12 months later. Typhoid Recommended for all volunteers. Generally given in oral form. Yellow Fever Required by the Malawian Ministry of Health for all volunteers entering Malawi. This involves a cost. Hepatitis B Recommended for all volunteers.

Rabies Recommended for all volunteers. This involves a cost. MMR (measles, mumps, rubella) Two doses recommended for all volunteers born after 1956.

Tetanus – diphtheria Recommended for all volunteers who have not received a tetanus - diphtheria immunisation within the last ten years.

Anti-malarial tablets: A malaria prophylaxis is essential for all volunteers. Please consult a medical practitioner before starting any of these medicines. **There are three main types:**

Drug Dosage Side Effects Note Lariam (Mefloquine) Once a week, starting 2 – 3 weeks before arrival (change if side effects are noticed) and continuing for 4 weeks after departure. Nausea, vomiting, dizziness, insomnia, nightmares. On some occasions depression, anxiety, hallucinations. Cheapest. Adverse side effects – not suitable for everyone. Please consult a medical practitioner before starting this medicine. Doxycycline Once daily, starting 2 days before arrival and continuing for 4 weeks after departure. Photosensitivity, exaggerated sunburn reactions, indigestion, nausea if taken on an empty stomach. Mid-range price. Recommended for longterm volunteers. Malarone Once daily, starting 2 days before arrival and continuing for 7 days after departure. Typically mild. On occasions abdominal pain, nausea, vomiting, headaches. Most expensive. Least side effects. Recommended for shortterm volunteers.

Medication: To avoid malaria, insect repellent with maximum 50% DEET should be bought before leaving for Malawi. Long-term volunteers may at times not have access to medical care whilst travelling and are advised to purchase a medical pack with materials for emergency self-treatment. Travellers' diarrhoea is very common, although most cases are mild and do not require antibiotics or anti-diarrhoeal drugs. In such a situation adequate fluid intake is normally sufficient, however volunteers may wish to purchase relevant medication before entering the country. HIV/AIDS is extremely prevalent in most parts of Malawi, and all volunteers are recommended to bring condoms before their arrival. We do however recommend all volunteers to carry their own personal first aid kit with the following items: (1)Antibiotics – a broad spectrum antibiotic such as a penicillin for general infections and either metronidazole or ciprofloxacin for bacterial diarrhoea (2) Antiseptic powder/ointment – cuts and sores get easily infected in a humid environment (3)Anti-diarrhoeal medication e.g. Imodium (4)Oral rehydration salts (5)Painkillers and anti-inflammatories e.g. Paracetamol, Ibuprofen (6)Antihistamines – for allergic reactions, Hydrocortisone cream – for allergic skin reactions or itchy bites: Bandages, plasters, gauze and adhesive tape , Syringes and sterile needles ,Thermometer ,Insect repellent with max 50% DEET ,Suntan lotion ,Antiseptic hand lotion , Please don't forget any of your usual medication such as the contraceptive pill, epi-pens etc. There are emergency first aid kits for travellers available at many stores or online which contain some or all of the above equipment in one pack.

During Your Stay

Volunteer Precautions: A – Malaria Malaria is a problem in Ruarwe and we strongly advise all volunteers to follow recommended precautions:

1. Make sure to take anti-malarial tablets for the recommended time before entering the country and after leaving.
2. Make sure to take your pills everyday around the same time and after eating (or every week if you are taking Larium).
3. Use insect repellent. Try to cover yourself in the evening, minimising as much bare skin as possible. Always use a mosquito net, and check to make sure any holes are covered.
4. HIV/AIDS :Volunteers are not at risk unless they have unprotected sex or receive injections or blood transfusions. Always use a condom. There are clinics in Chikwawa where volunteers can get free HIV tests and same-day results.
5. Bilharzia: Only swim in areas of the lake where the water is moving and clear . Avoid stagnant areas where there are many reeds. Make sure to get medication (Praziquantel) before leaving Malawi (these should be taken as a single dose with a large meal and plenty of water before bed approximately 6-8 weeks after leaving).
6. Giardia : Do not drink tap water unless it has been boiled, filtered or chemically disinfected. Do not drink unbottled beverages or drinks with ice. Do not eat fruits or vegetables unless they have been peeled or cooked. Avoid cooked foods that have been sitting at room temperature for a long period of time. Do not eat raw or undercooked meat or fish.

DIN Malawi Precautions: Volunteers that are worried about having contracted any of the above should report immediately to the project coordinator. Cases of severe illness are very rare but often occur due to lack of action.

In case a situation arises where a volunteer contracts malaria, Coartem or an equivalent is stocked. Coartem is one of the most successful and side-effect-free malaria treatments around and is suitable for use in most cases.

If the situation arises where a volunteer experiences sickness that is most likely not related to malaria, they will be referred to one of the following:

In all medical situations the volunteer is expected to cover the costs for their treatment, appointments, and travel to and from the clinic/hospital.

After Your Visit

Precautions: After volunteering with us at DIN Malawi recommends you take the following precautions before embarking on your journey home:

- i. Anti-malarial tablets should be taken for at least one week after leaving the country.
- ii. Tablets for treating bilharzia should be bought before leaving Malawi. These can be obtained from any pharmacy and the chemist will provide you with further information on how and when to take the tablets (you need to know your approximate weight in kg).
- iii. Due to the nature of the high HIV/AIDS rate in Malawi, DIN Malawi recommends volunteers to get tested once they are back in their home country.
- iv. If any volunteers plan to travel around Malawi or other parts of Africa once they have completed their work with the DIN Malawi, then they are strongly recommended to consult their GP to seek further advice on precautions for cholera, yellow fever etc.

We hope that all volunteers have a wonderful stay in Malawi with DIN, and we will do our very best to ensure the safety and well being of all our helpers.

If you need any more information about precautions to take before volunteering with us then please contact our Project Manager: dinorg2012@gmail.com